



Clustering of Dusty Star-Forming Galaxies

marco viero






Chary & Elbaz (2001)




UV from young, hot stars




Chary & Elbaz (2001)






Chary & Elbaz (2001)






Chary & Elbaz (2001)






Dole et al. 2006

Extragalactic Background Light (EBL)




Negative K-correction




Negative K-correction

get this data @ <http://www.ias.u-psud.fr/irgalaxies/model.php#Counts>




Béthermin et al. (2011)
arXiv:1010.1150

Redshift Distribution



Béthermin et al. (2011)
arXiv:1010.1150


Total Infrared Background



B  thermin et al. (2011)
arXiv:1010.1150

250 350 500

Total Infrared Background




Infrared Background
at $\lambda > 1$ mm (< 300 GHz)
dominated by
high-redshift sources

Béthermin et al. (2011)
arXiv:1010.1150

250 350 500

Total Infrared Background




Infrared Background
at $\lambda > 1$ mm (< 300 GHz)
dominated by
high-redshift sources

CMB bands at
220 GHz (1.4 mm)
148 GHz (2 mm)

Béthermin et al. (2011)
arXiv:1010.1150


250 350 500

Total Infrared Background




Béthermin et al. (2011)
 arXiv:1010.1150

Star Formation History



Le Floc'h et al. (2005)


What are they?



Le Floc'h et al. (2005)


What are they?

250 μm : flux density vs. redshift



What are they?

250 μ m: flux density vs. redshift



Typical **LIRGS** at $z \sim 1-3$
have flux densities
 $S_{250} \sim 1-10$ mJy

What are they?

Goal:

Measure the spatial distribution (clustering) of galaxies responsible for most of the SF in the Universes History

Goal:

Measure the spatial distribution (clustering) of galaxies responsible for most of the SF in the Universes History

1. Provides link to descendants/progenitors

Goal:


Measure the spatial distribution (clustering) of galaxies responsible for most of the SF in the Universes History

1. Provides link to descendants/progenitors
2. Informs galaxy formation models

Observing in the submillimeter




ZENITH - TRANSMISSION OF THE ATMOSPHERE IN DEPENDENCE OF THE PRECIPITABLE WATER VAPOR



Submm Observations

ZENITH - TRANSMISSION OF THE ATMOSPHERE IN DEPENDENCE OF THE PRECIPITABLE WATER VAPOR



MAMBO


AzTEC

SCUBA

SHARC

Submm Observations

ZENITH - TRANSMISSION OF THE ATMOSPHERE IN DEPENDENCE OF THE PRECIPITABLE WATER VAPOR






500



350



250




Submm Observations













SPIRE-like spider-web bolometer detector arrays...





SPIRE-like
spider-web bolometer
detector arrays...

... but a 1.8 m primary
resulting in beams:

36" - 250 μm

45" - 350 μm


60" - 500 μm

i.e., twice the size of SPIRE's
beams







BLAST GOODS-south



- Noise BGS-wide:
36 mJy beam⁻¹
21 mJy beam⁻¹
20 mJy beam⁻¹

- Noise BGS-deep:
11 mJy beam⁻¹
9 mJy beam⁻¹
6 mJy beam⁻¹

BLAST GOODS-south



- Noise BGS-wide:
36 mJy beam⁻¹
21 mJy beam⁻¹
20 mJy beam⁻¹

- Noise BGS-deep:
11 mJy beam⁻¹
9 mJy beam⁻¹
6 mJy beam⁻¹




image: J. Vieira

submm surveys





image: J. Vieira


submm surveys by year



Béthermin et al. (2011)
arXiv:1010.1150

Spitzer (Frayser 2009)
Herschel (Nguyen
2010)
APEX
CSO
CCAT


Confusion



U-band
 ~ 60 sources/
 arcmin^2

$\longleftrightarrow 10'$ U-band : MUSYC $\longrightarrow$

Confusion




U-band
 ~ 60 sources/
 arcmin^2

K-band
 ~ 45 sources/
 arcmin^2

$\longleftrightarrow 10' \xrightarrow{\text{K-band : MUSYC}}$

Confusion



Gaelen Marsden, UBC

- U-band
 ~ 60 sources/
 arcmin^2
- K-band
 ~ 45 sources/
 arcmin^2
- SPIRE
 ~ 0.5 sources/
 arcmin^2

← 10' SPIRE 250 μm →



Confusion

250 μm


$S > 20 \text{ mJy} : 1,200/\text{deg}^2$

$S < 20 \text{ mJy} : 480,000/\text{deg}^2$


Confusion



Confusion





50 brightest
(top 0.5%)




50 brightest
recovered

Confusion



70% of top
0.5%
recovered


Confusion




Model – Béthermin et al. (2011)

arXiv:1010.1150

Measured Number Counts




Measured Number Counts



Measured Number Counts


How do you measure
clustering when most of
what you see are
fluctuations?



$$P_{\text{shot}} = S^2 \frac{dN}{dS}$$


Poisson (shot) Noise

Star Forming Galaxies are
biased tracers of Dark Matter



Star Forming Galaxies are
biased tracers of Dark Matter


BLAST Power Spectrum



Viero et al. (2009)

arXiv: 0904.1200

BLAST Power Spectrum




Shot
Noise

Viero et al. (2009)

arXiv: 0904.1200

BLAST Power Spectrum



Shot
Noise


Clustering

Excess
Signal from
Galaxy
Clustering

Viero et al. (2009)
arXiv: 0904.1200

BLAST Power Spectrum

Clustering Signal
After Removal of
Shot Noise





Viero et al. (2009)

arXiv: 0904.1200



BLAST Power Spectrum

BLAST simulation...




BLAST Power Spectrum

BLAST simulation...



... Herschel observation

BLAST Power Spectrum




Planck Collaboration et al. (2011)

arXiv: 1101.2028


Planck Power Law

$$P(\ell) = P_{\text{shot}} + A (\ell / 1000)^{-n}$$




Addison et al. (in prep.)

Frequency-dependence of Power-law




Latest Power Spectra



see also Shirokoff et al. (2010) – arXiv: 1012.4788

Millea et al. (2011) – arXiv: 1102.5195

Latest Power Spectra




Viero et al. (2009)

arXiv: 0904.1200

$$P(k) = b^2 P_{\text{DM}}(k, z)$$

Linear Bias




$$P(k) = b^2 P_{\text{DM}}(k, z)$$

Linear Bias

see e.g., Cooray & Sheth (2000),
Zehavi et al. (2005, 2008)



see e.g., Cooray & Sheth (2000),
Zehavi et al. (2005, 2008)

- Clustering Signal made up of two regimes



see e.g., Cooray & Sheth (2000),
Zehavi et al. (2005, 2008)



- Clustering Signal made up of two regimes
- 2-halo: Linear Regime (large scales)



BLAST Halo Model

see e.g., Cooray & Sheth (2000),
Zehavi et al. (2005, 2008)



- Clustering Signal made up of two regimes
 - 2-halo: Linear Regime (large scales)
 - 1-halo: Non-Linear Regime (small scales)



BLAST Halo Model

see e.g., Cooray & Sheth (2000),
Zehavi et al. (2005, 2008)

- Clustering Signal made up of two regimes
 - 2-halo: Linear Regime (large scales)
 - 1-halo: Non-Linear Regime (small scales)



BLAST Halo Model

What are you *actually* fitting?

$$P_{1h}(k, z) = \int_{\mathcal{M}} n_{\text{halo}}(M, z) [2N_{\text{cen}}(M)N_{\text{sat}}(M)u_{\text{DM}}(k, z|M) + N_{\text{sat}}^2(M)u_{\text{DM}}^2(k, z|M)] dM / n_{\text{gal}}^2(z),$$

$$P_{2h}(k, z) = P_{\text{DM}}(k, z) \left[\int_{\mathcal{M}} n_{\text{halo}}(M, z) N_{\text{gal}}(M, z) \times b(M, z) u_{\text{DM}}(k, z|M) dM \right]^2 / n_{\text{gal}}^2(z).$$

$$n_{\text{gal}}(z) = \int_{\mathcal{M}} n_{\text{halo}}(M, z) \left[1 + \left(\frac{M}{M_1} \right)^\alpha \right] dM$$

and for each $M_{\text{min}}\text{-}\alpha$ pair, M_1 fixed to agree to source model by requiring:

$$\int_0^\infty \frac{dN}{dS dz}(S, z) dS = n_{\text{gal}}(z) dV_c(z)$$

What are you
actually fitting?

$$M_{\min} \approx 3 \times 10^{11} M_{\text{sun}}$$

$$\alpha \approx 1.1$$


$$P_{1h}(k, z) = \int_{\mathcal{M}} n_{\text{halo}}(M, z) [2N_{\text{cen}}(M)N_{\text{sat}}(M)u_{\text{DM}}(k, z|M) + N_{\text{sat}}^2(M)u_{\text{DM}}^2(k, z|M)] dM / n_{\text{gal}}^2(z),$$

$$P_{2h}(k, z) = P_{\text{DM}}(k, z) \left[\int_{\mathcal{M}} n_{\text{halo}}(M, z) N_{\text{gal}}(M, z) \times b(M, z) u_{\text{DM}}(k, z|M) dM \right]^2 / n_{\text{gal}}^2(z).$$

$$n_{\text{gal}}(z) = \int_{\mathcal{M}} n_{\text{halo}}(M, z) \left[1 + \left(\frac{M}{M_1} \right)^\alpha \right] dM$$


and for each $M_{\min}$ - α pair, M_1 fixed to agree to source model by requiring:

$$\int_0^\infty \frac{dN}{dS dz}(S, z) dS = n_{\text{gal}}(z) dV_c(z)$$



Viero et al. (2009)

BLAST Halo Model




$$n_{\text{gal}}(z) = \int_{\mathcal{M}} n_{\text{halo}}$$

and for each $M_{\min}$ - α pair, M_1 fixed to agree to source model by requiring:

$$\int_0^{\infty} \frac{dN}{dS dz}(S, z) dS = n_{\text{gal}}(z) dV_c(z)$$

BLAST Halo Model


Halo Model of **Mattia Negrello** & Source Model of **Lagache** 2004



Viero et al. (2009)


arXiv: 0904.1200

BLAST Halo Model



Agrees
w/ Planck!

of Mattia Negrello &
et al of Lagache 2004



Viero et al. (2009)


arXiv: 0904.1200

BLAST Halo Model

At $z \sim 2$

- $b_{\text{linear}} \approx 4.0$
- $M_{\text{eff}} \approx 6 \times 10^{12} M_{\text{sun}}$
- $M_{\text{min}} \approx 3 \times 10^{11} M_{\text{sun}}$
- $b_{\text{eff}} \approx 2.4$

Halo Model of **Mattia Negrello** &
Source Model of **Lagache** 2004



Viero et al. (2009)

arXiv: 0904.1200

BLAST Halo Model

At $z \sim 2$

- $b_{\text{linear}} \approx 4.0$

- $M_{\text{eff}} \approx 6 \times 10^{12} M_{\text{sun}}$

- $M_{\text{min}} \approx 3 \times 10^{11} M_{\text{sun}}$

- $b_{\text{eff}} \approx 2.4$

Sub-millimetre galaxies reside in dark matter halos with masses greater than $3 \times 10^{11} M_{\odot}$

A. Amblard¹, A. Cooray^{1,2}, P. Serra¹, B. Altieri³, V. Arumugam⁴, H. Aussel⁵, A. Blain², J. Bock^{2,6}, A. Boselli⁷, V. Buat⁷, N. Castro-Rodríguez^{8,9}, A. Cava^{8,9}, P. Chanial¹⁰, E. Chapin¹¹, D.L. Clements¹⁰, A. Conley¹², L. Conversi³, C.D. Dowell^{2,6}, E. Dwek¹³, S. Eales¹⁴, D. Elbaz⁵, D. Farrah¹⁵, A. Franceschini¹⁶, W. Gear¹⁴, J. Glenn¹², M. Griffin¹⁴, M. Halpern¹¹, E. Hatziminaoglou¹⁷, E. Ibar¹⁸, K. Isaak¹⁴, R.J. Ivison^{18,4}, A.A. Khostovan¹, G. Lagache¹⁹, L. Levenson^{2,6}, N. Lu^{2,20}, S. Madden⁵, B. Maffei²¹, G. Mainetti¹⁶, L. Marchetti¹⁶, G. Marsden¹¹, K. Mitchell-Wynne¹, H.T. Nguyen^{6,2}, B. O'Halloran¹⁰, S.J. Oliver¹⁵, A. Omont²², M.J. Page²³, P. Panuzzo⁵, A. Papageorgiou¹⁴, C.P. Pearson^{23,24}, I. Pérez-Fournon^{8,9}, M. Pohlen¹⁴, N. Rangwala¹², I.G. Roseboom¹⁵, M. Rowan-Robinson¹⁰, M. Sánchez Portal¹, M. Sygda¹, L. Vigotti¹

Using the halo model fits, we estimate the minimum dark matter mass scale for dusty star-forming galaxies at the peak of the star formation history of the universe to be $\log_{10} M_{\text{min}}/M_{\odot} = 11.5^{+0.7}_{-0.2}$ at 350 μm with a bias factor for the galaxies of $2.4^{+1.0}_{-0.2}$. The minimum halo masses $\log_{10} M_{\text{min}}/M_{\odot}$ at 250 and 500 μm are $11.1^{+1.0}_{-0.6}$ and $11.8^{+0.4}_{-0.3}$, respectively. The corresponding bias factors for the galaxies are $2.0^{+0.9}_{-0.1}$ and $2.8^{+0.4}_{-0.5}$ at 250 and 500 μm , respectively. The differences in the minimum halo masses and the bias factors between the three wavelengths are likely due a combination of effects including overall calibration uncertainties, the fact that at longer wave-

arXiv: 0904.1200

Confirmed
by Herschel!

BLAST Halo Model

At $z \sim 2$

- $b_{\text{linear}} \approx 4.0$

- $M_{\text{eff}} \approx 6 \times 10^{12} M_{\text{sun}}$

- $M_{\text{min}} \approx 3 \times 10^{11} M_{\text{sun}}$

- $b_{\text{eff}} \approx 2.4$


Sub-millimetre galaxies reside in dark matter halos with masses greater than $3 \times 10^{11} M_{\odot}$

A. Amblard¹, A. Cooray^{1,2}, P. Serra¹, B. Altieri³, V. Arumugam⁴, H. Aussel⁵, A. Blain², J. Bock^{2,6}, A. Boselli⁷, V. Buat⁷, N. Castro-Rodríguez^{8,9}, A. Cava^{8,9}, P. Chanial¹⁰, E. Chapin¹¹, D.L. Clements¹⁰, A. Conley¹², L. Conversi³, C.D. Dowell^{2,6}, E. Dwek¹³, S. Eales¹⁴, D. Elbaz⁵, D. Farrah¹⁵, A. Franceschini¹⁶, W. Gear¹⁴, J. Glenn¹², M. Griffin¹⁴, M. Halpern¹¹, E. Hatziminaoglou¹⁷, E. Ibar¹⁸, K. Isaak¹⁴, R.J. Ivison^{18,4}, A.A. Khostovan¹, G. Lagache¹⁹, L. Levenson^{2,6}, N. Lu^{2,20}, S. Madden⁵, B. Maffei²¹, G. Mainetti¹⁶, L. Marchetti¹⁶, G. Marsden¹¹, K. Mitchell-Wynne¹, H.T. Nguyen^{6,2}, B. O'Halloran¹⁰, S.J. Oliver¹⁵, A. Omont²², M.J. Page²³, P. Panuzzo⁵, A. Papageorgiou¹⁴, C.P. Pearson^{23,24}, I. Pérez-Fournon^{8,9}, M. Pohlen¹⁴, N. Rangwala¹², I.G. Roseboom¹⁵, M. Rowan-Robinson¹⁰, M. Sánchez Portal¹, M. Sygga¹, L. Vigor¹

Using the halo model fits, we estimate the minimum dark matter mass scale for dusty star-forming galaxies at the peak of the star formation history of the universe to be $\log_{10} M_{\text{min}}/M_{\odot} = 11.5^{+0.7}_{-0.2}$ at 350 μm with a bias factor for the galaxies of $2.4^{+1.0}_{-0.2}$. The minimum halo masses $\log_{10} M_{\text{min}}/M_{\odot}$ at 250 and 500 μm are $11.1^{+1.0}_{-0.6}$ and $11.8^{+0.4}_{-0.3}$, respectively. The corresponding bias factors for the galaxies are $2.0^{+0.9}_{-0.1}$ and $2.8^{+0.4}_{-0.5}$ at 250 and 500 μm , respectively. The differences in the minimum halo masses and the bias factors between the three wavelengths are likely due a combination of effects including overall calibration uncertainties, the fact that at longer wave-

Confirmed
by Herschel!

BLAST Halo Model




- Fix $M_1/M_{\min}=10$
- Use B10 model for Poisson term

Planck Collaboration et al. (2011)

arXiv: 1101.2028

Planck Halo Model




- Fix $M_1/M_{\min}=10$
- Use B10 model for Poisson term

Planck Collaboration et al. (2011)

arXiv: 1101.2028


Planck Halo Model



Amblard et al. (2011)

arXiv: 1101.1080


Herschel Halo Model




Amblard et al. (2011)

arXiv: 1101.1080

Herschel Halo Model


The background of the slide is a Cosmic Microwave Background (CMB) fluctuation map. It shows a complex pattern of temperature variations across the sky, represented by a color scale from blue (cooler) to yellow and red (warmer). The pattern consists of numerous small, irregular patches of different colors, creating a grainy, textured appearance. The text 'cmb?' is centered over this map.

cmb?



The high- ℓ CMB sky


CMB



The high- ℓ CMB sky

CMB

Galaxies




The high- ℓ CMB sky


CMB

Galaxies

SZ Clusters




The high- ℓ CMB sky




Dunkley et al. 1009.0866

CMB Foregrounds




Dunkley et al. 1009.0866

CMB Foregrounds




Dunkley et al. 1009.0866

CMB Foregrounds




Dunkley et al. 1009.0866

CMB Foregrounds



Dunkley et al. 1009.0866

CMB Foregrounds



Dunkley et al. 1009.0866


CMB Foregrounds

Atacama Cosmology Telescope (ACT)



ACT Southern Field

Hajian et al (2010)



- 300 deg²
- 1' resolution
- 3 bands


148 GHz (~2.0 mm)

218 GHz (~1.4 mm)

277 GHz (~1.1 mm)





$\sim 9 \text{ deg}^2$




Noise r.m.s.:
36.7 mJy beam⁻¹
27.2 mJy beam⁻¹
19.1 mJy beam⁻¹

Overlap: BLAST-SEP Field

ACT 148 GHz (2 mm)




IRAS 100 μm




Mean Flux $\sim 1.1 \text{ MJy/sr}$

BLAST-Maps made with **SANEPIC**



BLAST-Maps made with **SANEPIC**





Cross-Correlations



Isolate Dusty

Galaxies from the

Rest




Cross-Correlations *Isolate* Dusty Galaxies from the Rest




ACT X BLAST

Cross-correlation in Fourier Space




Cross-correlation in Fourier Space




Fourier Domain Filter


Cross-Correlate BLAST and ACT




Hajian & Viero et al.
arXiv:1101.1517




Hajian & Viero et al.
arXiv:1101.1517



Hajian & Viero et al.
 arXiv:1101.1517




Hajian & Viero et al.
arXiv:1101.1517



Hajian & Viero et al.
arXiv:1101.1517

Fit a Linear bias **simultaneously** to each band

Source model of [Béthermin et al. \(2010\)](#)
& Linear bias model of [Graeme Addison](#)



Constrains the evolution using redshift information


Finds bias strongly dependent on model

[Hajian & Viero et al. \(2011\)](#)

[arXiv: 1101.1517](#)

Fit a Linear bias **simultaneously** to each band

Source model of [Béthermin et al. \(2010\)](#)
& Linear bias model of [Graeme Addison](#)




Constrains the evolution using redshift information

Finds bias strongly dependent on model

[Hajian & Viero et al. \(2011\)](#)

[arXiv: 1101.1517](#)

ACT X BLAST



Model: Marsden et al. 1010.1176

Model: Béthermin et al. 1010.1150

- Auto-Frequency Correlations:

- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs

- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs
 - ✓ Constrain physical properties of Host Halos, such as bias and $M_{\min}$

- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs
 - ✓ Constrain physical properties of Host Halos, such as bias and $M_{\min}$
- Cross-Frequency Correlations:

- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs
 - ✓ Constrain physical properties of Host Halos, such as bias and M_{min}
- Cross-Frequency Correlations:
 - ✓ Isolate Dusty Galaxies in CMB maps

- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs
 - ✓ Constrain physical properties of Host Halos, such as bias and $M_{\min}$
- Cross-Frequency Correlations:
 - ✓ Isolate Dusty Galaxies in CMB maps
 - ✓ Identify the High- z Clustering Signal


- Auto-Frequency Correlations:
 - ✓ Measure *Clustering* of DSFGs
 - ✓ Constrain physical properties of Host Halos, such as bias and $M_{\min}$
- Cross-Frequency Correlations:
 - ✓ Isolate Dusty Galaxies in CMB maps
 - ✓ Identify the High- z Clustering Signal
 - ✓ Constrain Phenomenological Galaxy Models

future surveys



image: Joaquin Vieira

Neutral



Ionized

Zahn et al. (2005)


At $z < 30$,
Reionized
bubbles form
around UV
sources

Bubbles Moving
with respect to
CMB, thus
distort it

Strength and
Shape of kSZ
signal depends
on details of
Reionization

Reionization

Again, Problem of Foregrounds




Courtesy Joaquin Vieira

marco viero




53

To Get Weak kSZ Signal, you need **SPIRE**!



**Pesky foreground we
want to remove**





**Reionization signal we
want to measure**

Measuring Epoch of Reionization

Herschel/SPIRE X SPT-Deep Field

SPT 100 deg² deep field is the deepest mm map in existence and will remain so for the next decade.



Courtesy Joaquin Vieira

marco viero





55

Measuring Epoch of Reionization

Herschel/SPIRE X SPT-Deep Field

SPT 100 deg² deep field is the deepest mm map in existence and will remain so for the next decade.



Given 79 hours to
map a 100 deg² with
SPIRE

Measuring Epoch of Reionization

Herschel/SPIRE X SPT-Deep Field

SPT 100 deg² deep field is
the deepest mm map in
existence and will remain so
for the next decade



The HOTAC called
this proposal a
"Must-Do"

Given 79 hours to
map a 100 deg² with
SPIRE



Courtesy Joaquin Vieira

marco viero



55

end

