


**Ay1 – Lecture 14**

# **Galaxy Formation and Evolution**

Abell 2744 (HST)





# 14.1 Galaxy Evolution: the Basics



# Galaxies Must Evolve

- Stars evolve: they are born from ISM, evolve, shed envelopes or explode, enriching the ISM, more stars are born...
- Structure evolves: density fluctuations collapse and merge in a hierarchical fashion

## Galaxy Evolution

Assembly of the mass



Conversion of gas into stars and v.v., feedback processes

DM dominated

Cannot be observed directly,  
but may be inferred

Easy to model, mainly  
dissipationless

This is what is observed, and  
where energy is generated  
Dissipative, and very hard to  
model

# Evolution Timescales and Evidence

**Timescales for galactic evolution** span wide range:

~ 100 Myr - galaxy free-fall and cooling time scales

10 -100 Myr - lifetimes of massive stars

10 -100 Myr - lifetime of the bright phase of a luminous  
Active Galactic Nucleus (?)

Few  $\times$  100 Myr - rotation period of spiral galaxy

~ Gyr - time required for two galaxies to merge

~ 10 Gyr - age of the Universe


**Observational evidence** for evolution is found in:

- Stellar populations in the Milky Way (e.g., metallicity as a function of stellar age, etc.)
- Systematics of nearby galaxy properties
- **Properties of distant galaxies seen at earlier epoch**




# Theoretical Tools and Approaches

- 1. Assembly of the mass:** numerical modeling of structure formation. Fairly well advanced, but it is hard to treat any dissipative processes very accurately. Well constrained from large-scale structure formation
- 2. Evolution of stellar populations:** based on stellar evolution models, and fairly well understood. Lots of parameters: the stellar initial mass function, star formation history, stellar evolutionary tracks and spectra as functions of metallicity. Poorly constrained






# A Group Formation Simulation by A. Kravtsov





# Snapshots from the Virgo Consortium's "Millenium" simulation

125 Mpc/h



**Credit:**

**Volker Springel**



**Z = 18.3**

31.25 Mpc/h

A visualization of the cosmic web at redshift Z = 18.3. The image shows a complex network of filaments and nodes, rendered in shades of purple and blue. A horizontal scale bar is positioned in the center, labeled "31.25 Mpc/h". The overall structure is highly interconnected and filamentary.



$Z = 5.7$


31.25 Mpc/h

A visualization of the cosmic web at redshift  $Z = 5.7$ . The image shows a complex network of filaments and nodes, with colors ranging from dark purple to bright yellow. A scale bar in the center indicates a distance of 31.25 Mpc/h. The filaments are interconnected, forming a dense web of structure. The nodes are the points where filaments intersect, often appearing as brighter yellow or white spots. The overall appearance is that of a highly interconnected, filamentary structure.




$Z = 1.4$

31.25 Mpc/h









**$z=1.7$**

# **The Milky Way Formation Simulation by Diemand et al. (Dark Matter Only)**






# Stellar Population Synthesis Models

We can synthesize predicted galaxy spectra as a function of time by assuming the following:

- Star formation rate as a function of time
- Initial mass function of stars
- Libraries of stellar spectra for stars of different masses, metallicities and ages, etc.
- Stellar evolutionary tracks (isochrones)






# Modeling Evolution of Stellar Pop's


- Stellar evolution is relatively well understood both observationally and theoretically; the key points:
  - Massive stars are very hot, blue, very luminous, and have very short lives; they dominate the restframe UV light
  - Thus we expect largest effects in the bluer parts of the spectrum
- Star formation histories are a key assumption:
  - Ellipticals are best fit by a burst of early star formation followed by a relatively “passive evolution” where they fade and get redder quickly
  - Spirals are best fit by a slowly declining or nearly constant star formation – they stay bluer and don't fade as much

# Predicted Spectral Evolution




Generally, models fade in time, and faster in the UV than in the IR

# Colors of Stellar Populations: Differences in Star Formation Histories





## 14.2 Observations of Galaxy Evolution



# Observational Tools and Approaches

- **Deep imaging surveys** and source counts, at wavelengths from UV to FIR
  - Sources are always selected in emission, and any given band has its own selection effects and other peculiarities
  - With enough bandpasses, one can estimate “photometric redshifts”, essentially very low resolution spectroscopy; may be unreliable
  - Measurements of galaxy clustering provide additional information
- **Deep spectroscopic redshift surveys:** redshifts are usually obtained in the visible, regardless of how the sources are selected
  - As a bonus, one can also estimate current star formation rates and rough chemical abundances from the spectra
- **Diffuse extragalactic backgrounds:** an integrated emission from all sources, regardless of the flux or surface brightness limits
  - Extremely hard to do
  - No redshift information




# Observing Galaxy Evolution

- If redshifts are not available, we can do source counts as a function of limiting flux or magnitude; and colors as a function of magnitude (acting as a proxy for distance - not a great approximation)
- But you really do need redshifts, to get a true evolution in time, and disentangle the various evolution effects
- The field is split observationally:
  - **Unobscured star formation** evolution: most of the energy emerging in the restframe UV, observed in the visible/NIR
  - **Obscured star formation**: energy from young stars reprocessed by dust to emerge in FIR/sub-mm
  - They have different limitations and selection effects

# Source Counts: The Effect of Evolution

$\log N$  (per unit area  
and unit flux or mag)

(at a fixed cosmology!)



Evolution

No evolution

**Luminosity evolution**  
moves fainter sources (more  
distant and more numerous) to brighter  
fluxes, thus producing excess counts, since  
generally galaxies were brighter in the past

**Density evolution** means that there was some galaxy  
merging, so there were more fainter pieces in the past,  
thus also producing excess counts at the faint end

In order to distinguish between  
the two evolution mechanisms,  
redshifts are necessary




# Galaxy Counts in Practice

The deepest galaxy counts to date come from HST deep and ultra-deep observations, reaching down to  $\sim 29^{\text{th}}$  mag

All show excess over the no-evolution models, and more in the bluer bands

The extrapolated total count is  $\sim 10^{11}$  galaxies over the entire sky



A proven powerful combination is to use deep HST imaging (e.g., HDF N and S, HUDF, GOODS field, etc.) and Keck or other 6 to 10-m class telescope for spectroscopy.


Various deep fields also have multi-wavelength data from Chandra, VLA, Spitzer ...






# Deep Redshift Surveys

- To really understand what is going on, separate the effects of luminosity and density evolution, and break the degeneracy between distance and intrinsic luminosity at a given flux, we need redshifts
- To go beyond  $z > 1$ , we have to go faint, e.g. to  $R > 23 - 24$  mag



# Evolution of Galaxy Sizes

HST imaging suggests that galaxies were smaller in the past







# Evolution of the Merger Rate

*Merger fraction*

$$\sim (1+z)^{3.4}$$




(LeFevre et al. 2000)

Good evidence for a rapid rise in merging fraction at higher  $z$ 's, but conversion to mass assembly rate is not straightforward

# Scaling Relations as Evolution Probes

- Studies of the Fundamental Plane out to  $z > 1$  indicate that ellipticals were brighter in the past, but the data are consistent with a model where they are formed at high redshifts ( $z > 3$ , say) and evolve nearly passively since then



- Data on galaxy colors and line strengths are also consistent with that picture
- Studies of the *Tully-Fisher relation* at high  $z$ 's are much less conclusive



# Galaxies in the past were...

- **Bluer**
- **Brighter at a given mass**  
**but**
- **Smaller mass on the average**
- **Smaller in size**
- **More numerous**



# 14.3 The Cosmic Star Formation History







# M82, a Prototypical Starburst Galaxy




The spectrum of M82, UV to sub-mm



Unobscured  
starlight

Reprocessed  
radiation from  
dust



Distant Galaxy in the Hubble Ultra Deep Field

Spitzer Space Telescope • IRAC  
Hubble Space Telescope • ACS • NICMOS




# Cosmic Star Formation History

From various luminosity densities converted to star formation rates, we can construct a possible history of the comoving SFR density

At face value it implies the universe was much more active in the past ( $z \sim 1 - 2$ ) but what happens earlier is unclear

There are many complications of interpretation, including the reliability of each SFR diagnostic, dust extinction, incompleteness, etc.




# Now pushing to $z \sim 6$ (and beyond?)

Use the color dropout technique to identify high- $z$  galaxy candidates in deep HST images: different color bins give different redshift shells. Then add up the light.

There seems to be a rollover at  $z > 5 - 6$ : the epoch of the initial galaxy build-up?


*(Bouwens & Illingworth 2006)*






# Build-up of Stellar Mass Density

Co-moving stellar mass density grew rapidly from  $z \sim 3$  to  $z \sim 1$ , but has not changed much since then



# Build-up of Stellar Mass Density

... and the trend seems to continue out to  $z \sim 6$  (and beyond?)






# All Starlight in the Universe

- Any deep survey is limited in flux and surface brightness: some fainter and/or more diffuse sources are likely missed; thus, our source counts give us only a lower limit to the total energy emitted by evolving galaxies
- An alternative approach is to measure *integrated diffuse backgrounds, due to all sources*
  - This is really hard to do, for many reasons
  - Redshifts are lost, but at least the energy census is complete
- The total energy in the diffuse extragalactic backgrounds from UV to sub-mm is  $\sim 100 \text{ nW m}^{-2} \text{ sr}^{-1}$  ( $\pm 50\%$  or so)
  - This is distributed roughly equally between the UV/Opt (unobscured SF) and FIR/sub-mm (obscured SF)
  - A few percent of the total is contributed by AGN
  - This is only a few percent of the CMB

# Diffuse Optical and IR Backgrounds



Unobscured component  
(restframe UV, obs. Optical/NIR)

Obscured component  
(restframe FIR, obs. FIR/sub-mm)





**14.4 Cosmic Web:  
The Intergalactic Medium**


# Intergalactic Medium (IGM)

- Essentially, baryons between galaxies
- Its density evolution follows the large scale structure formation, and the potential wells defined by the dark matter, forming a web of filaments, the co-called “**Cosmic Web**”
- Gas falls into galaxies, where it serves as a replenishment fuel for star formation
- Likewise, enriched gas is driven from galaxies through the radiatively and SN powered **galactic winds**, which chemically enriches the IGM
- Chemical evolution of galaxies and IGM thus track each other
- Star formation and active galactic nuclei provide the **ionizing flux** for the IGM



# Cosmic Web: Numerical Simulations


Our lines of sight towards some luminous background sources intersect a range of gas densities, condensed clouds, galaxies ...



*(from R. Cen)*

# QSO Absorption Line Systems

- An alternative to searching for galaxies by their *emission* properties is to search for them by their *absorption*
- Quasars are very luminous objects and have very blue colours which make them relatively easy to detect at high redshifts
- GRB are also used
- Note that this has *different selection effects* than the traditional imaging surveys: not by luminosity or surface brightness, but by the size and column density






# Types of QSO Absorption Lines

- **Lyman alpha forest:**
  - Numerous, weak lines from low-density hydrogen clouds
  - Proto-galactic clouds, with low density, they are not galaxies
- **Lyman Limit Systems (LLS) and “Damped” Lyman alpha (DLA) absorption lines:**
  - Rare, strong hydrogen absorption, high column densities
  - Coming from intervening galaxies
  - An intervening galaxies often produce both metal and damped Lyman alpha absorptions
- **Helium** equivalents are seen in the far UV part of the spectrum
- **“Metal” absorption lines**
  - Absorption lines from heavy elements, e.g., C, Si, Mg, Al, Fe
  - Most are from intervening galaxies

# Types of QSO Absorption Systems


PKS 0454+039  $z=1.34$






# A Damped Lyman $\alpha$ System

Q1331+170  $z_{\text{em}}=2.084$   $z_{\text{abs}}=1.7764$  (WHT)



# Evolution of the Hydrogen Absorbers







# Galactic Winds

Starburst can drive winds of enriched gas (e.g., from supernova ejecta) out to the intergalactic medium. This gas can then be accreted again by galaxies. In a disk galaxy, the winds are generally bipolar outflows


M82 (CXO): X-ray



M82 (Subaru): H $\alpha$  + optical



Numerical Simulation





## 14.5 The Earliest Galaxies

A visualization of the cosmic web, showing a dense network of blue filaments and nodes against a black background. Numerous bright yellow and orange points are scattered throughout, representing early galaxies. The text "14.5 The Earliest Galaxies" is overlaid in yellow in the upper left quadrant.

# Galaxy Formation

- The early stages of galaxy evolution - but there is no clear-cut boundary, and it also has two principal aspects: assembly of the mass, and conversion of gas into stars
- Must be related to large-scale (hierarchical) structure formation, plus the dissipative processes
- Probably closely related to the formation of the massive central black holes as well
- Generally, we think of massive galaxy formation at high redshifts ( $z \sim 3 - 10$ , say); dwarfs may be still forming now
- Observations have found populations of what must be young galaxies (ages  $< 1$  Gyr), ostensibly progenitors of large galaxies today, at  $z \sim 5 - 7$
- The frontier is now at  $z \sim 7 - 20$ , the so-called Reionization Era




# A General Outline

- The smallest scale density fluctuations keep collapsing, with baryons falling into the potential wells dominated by the dark matter, achieving high densities through cooling
  - This process starts right after the recombination at  $z \sim 1100$
- Once the gas densities are high enough, star formation ignites
  - This probably happens around  $z \sim 20 - 30$
  - By  $z \sim 6$ , UV radiation from young galaxies reionizes the universe
- These protogalactic fragments keep merging, forming larger objects in a hierarchical fashion ever since then
- Star formation enriches the gas, and some of it is expelled in the intergalactic medium, while more gas keeps falling in
- If a central massive black hole forms, the energy release from it can also create a considerable feedback on the young host galaxy

# An Outline of the Early Cosmic History

*(illustration from Avi Loeb)*



↑  
**Recombination:**  
Release of the  
CMBR

↑  
**Dark Ages:**  
Collapse of  
Density  
Fluctuations

↑  
**Reionization  
Era:**  
The Cosmic  
Renaissance

↑  
Galaxy  
evolution  
begins




# Energy Release From Forming Galaxies

- Release of the binding energy from a collapsing protogalaxy  
Typically  $\sim 10^{59}$  erg
- Thermonuclear burning in stars  
Typically  $\sim 10^{60}$  erg
- Release of the binding energy from the collapsing protostars  
Typically  $\sim 10^{59}$  erg
- Energy input from an active nucleus, if present  
Up to  $\sim 10^{60}$  erg





# Emission Line Search for Young Galaxies

Spectra of star forming regions have strong recombination lines of hydrogen and various ions, e.g., [O III]



The strongest of them is expected to be the hydrogen **Lyman  $\alpha$**  line (unless absorbed by gas or dust)




# Long-Slit Spectroscopy + Serendipity





# Narrow-Band Imaging

A greatly increased contrast for an object with a strong line emission





# A Galaxy at $z \sim 7$

## IOK-1 at $z = 6.96$



(Iye et al. 2006; Subaru)

Discovered using narrow band imaging technique



# The Lyman-Break Method

Absorption by the interstellar and intergalactic hydrogen of the UV flux blueward of the Ly alpha line, and especially the Lyman limit, creates a continuum break which is easily detectable by multicolor imaging





# Color-Selected Candidate High-z Galaxies


7 star-forming galaxies located  $8.5 < z < 12$

$5\sigma$  detections in (160W+140W+125W) stack ( $m_{AB} < 30.1$ )

$2\sigma$  rejection in ultra-deep F105W ( $m_{AB} > 31.0$ )

$2\sigma$  rejection in ACS BViz ( $m_{AB} > 31.3$ )

Ellis et al (2013) Ap J Lett 763, L7



$z=11.9?$  380 Myr

$z=9.5$  520 Myr

$z=9.5$  520 Myr


$z=8.8$  570 Myr

$z=8.8$  570 Myr

$z=8.6$  590 Myr

$z=8.6$  590 Myr


# Star formation density of LBGs



Monotonically declining population to  $z \sim 6$  and beyond

Drop of  $\times 8$  in UV luminosity density over  $2 < z < 6$

Bouwens et al (2009, 2011)



**14.6 Reionization Era:  
The Cosmic Renaissance**



# What is the Reionization Era?

A Schematic Outline of the Cosmic History

## The Cosmic Reionization Era

(The Cosmic Renaissance)

Time since the Big Bang (years)

~ 300 thousand

DM Halos Form

~ 500 million

Pop III Stars, Early BH


Pop II +OMR, SMBH  
~ 1 billion

Evolution & Growth

~ 9 billion

Pop I ...

~ 13 billion



←The Big Bang

The Universe filled with ionized gas

←The Universe becomes neutral and opaque

The Dark Ages start

Galaxies and Quasars begin to form  
The Reionization starts

The Cosmic Renaissance  
The Dark Ages end

←Reionization complete, the Universe becomes transparent again



Galaxies evolve

The Solar System forms

Today: Astronomers figure it all out!


# The First Stars

Gas infall into the potential wells of the dark matter fluctuations leads to increased density, formation of  $H_2$ , molecular line cooling, further condensation and cloud fragmentation, leading to the formation of the **first stars**



# Population III Stars: Hot and Luminous

They can easily reionize the universe by  $z \sim 6$






# Population III Supernovae

- Early enrichment of the protogalactic gas
- Transition to the “normal” Pop II star formation and IMF when the metallicity reaches a critical value  $Z_{\text{crit}} \sim 10^{-3.5} Z_{\odot}$

Simulated Pop III SN shell after  $\sim 10^6$  yr



Distrib. of metals (red)

*(from Bromm et al. 2003)*




# GRB 090423 at $z \sim 8.2$ (?)

The current record holder –  
no details available yet




“Gunn-Peterson like” troughs are now observed along all available lines-of-sight at  $z \sim 6$





# Looking Even Deeper: The 21cm Line

We can in principle image H I condensations in the still neutral, pre-ionization universe using the 21cm line. Several experiments are now being constructed or planned to do this, e.g., the Mileura Wide-Field Array in Australia, or the Square Kilometer Array (SKA)



*(Simulations of  $z = 8.5$  H I, from P. Madau)*