

The most important forbidden lines in the interstellar medium. Only CI, CII, OI, SiII, SII, and FeII are present in the neutral medium. They are also present in the ionised medium, but generally in smaller amounts than more ionised species. The collision strengths  $\Omega_{ul}$  are for collisions with electrons at a temperature of  $10^4$  K. The critical densities  $n_{crit} = A_{ul}/\langle\sigma_{ul}v\rangle$  correspond to collisions either with electrons (for  $Te \approx 10^4$  K), or with  $H_2$  molecules when between round brackets (for  $T_k \approx 100$  K).

Ion	Transition l-u	$\lambda$ $\mu\text{m}$	$A_{ul}$ $\text{s}^{-1}$	$\Omega_{ul}$	$n_{crit}$ $\text{cm}^{-3}$
C I	$^3P_0 - ^3P_1$	609.1354	$7.93 \times 10^{-8}$	–	(500)
	$^3P_1 - ^3P_2$	370.4151	$2.65 \times 10^{-7}$	–	(3000)
C II	$^2P_{1/2} - ^2P_{3/2}$	157.741	$2.4 \times 10^{-6}$	1.80	47 (3000)
N II	$^3P_0 - ^3P_1$	205.3	$2.07 \times 10^{-6}$	0.41	41
	$^3P_1 - ^3P_2$	121.889	$7.46 \times 10^{-6}$	1.38	256
	$^3P_2 - ^1D_2$	0.65834	$2.73 \times 10^{-3}$	2.99	7700
	$^3P_1 - ^1D_2$	0.65481	$9.20 \times 10^{-4}$	2.99	7700
N III	$^2P_{1/2} - ^2P_{3/2}$	57.317	$4.8 \times 10^{-5}$	1.2	1880
O I	$^3P_2 - ^3P_1$	63.184	$8.95 \times 10^{-5}$	–	$2.3 \times 10^4$ ( $5 \times 10^5$ )
	$^3P_1 - ^3P_0$	145.525	$1.7 \times 10^{-5}$	–	3400 ( $1 \times 10^5$ )
	$^3P_2 - ^1D_2$	0.63003	$6.3 \times 10^{-3}$	–	$1.8 \times 10^6$
O II	$^4S_{3/2} - ^2D_{5/2}$	0.37288	$3.6 \times 10^{-5}$	0.88	1160
	$^4S_{3/2} - ^2D_{3/2}$	0.37260	$1.8 \times 10^{-4}$	0.59	3890
O III	$^3P_0 - ^3P_1$	88.356	$2.62 \times 10^{-5}$	0.39	461
	$^3P_1 - ^3P_2$	51.815	$9.76 \times 10^{-5}$	0.95	3250
	$^3P_2 - ^1D_2$	0.50069	$1.81 \times 10^{-2}$	2.50	$6.4 \times 10^5$
	$^3P_1 - ^1D_2$	0.49589	$6.21 \times 10^{-3}$	2.50	$6.4 \times 10^5$
	$^1D_2 - ^1S_0$	0.43632	1.70	0.40	$2.4 \times 10^7$
Ne II	$^2P_{1/2} - ^2P_{3/2}$	12.8136	$8.6 \times 10^{-3}$	0.37	$5.9 \times 10^5$
Ne III	$^3P_2 - ^3P_1$	15.5551	$3.1 \times 10^{-2}$	0.60	$1.27 \times 10^5$
	$^3P_1 - ^3P_0$	36.0135	$5.2 \times 10^{-3}$	0.21	$1.82 \times 10^4$
Si II	$^2P_{1/2} - ^2P_{3/2}$	34.8152	$2.17 \times 10^{-4}$	7.7	$(3.4 \times 10^5)$
S II	$^4S_{3/2} - ^2D_{5/2}$	0.67164	$2.60 \times 10^{-4}$	4.7	1240
	$^4S_{3/2} - ^2D_{3/2}$	0.67308	$8.82 \times 10^{-4}$	3.1	3270
S III	$^3P_0 - ^3P_1$	33.4810	$4.72 \times 10^{-4}$	4.0	1780
	$^3P_1 - ^3P_2$	18.7130	$2.07 \times 10^{-3}$	7.9	$1.4 \times 10^4$
S IV	$^2P_{1/2} - ^2P_{3/2}$	10.5105	$7.1 \times 10^{-3}$	8.5	$5.0 \times 10^4$
Ar II	$^2P_{1/2} - ^2P_{3/2}$	6.9853	$5.3 \times 10^{-2}$	2.9	$1.72 \times 10^6$
Ar III	$^3P_2 - ^3P_1$	8.9914	$3.08 \times 10^{-2}$	3.1	$2.75 \times 10^5$
	$^3P_1 - ^3P_0$	21.8293	$5.17 \times 10^{-3}$	1.3	$3.0 \times 10^4$
Fe II	$^6D_{7/2} - ^6D_{5/2}$	35.3491	$1.57 \times 10^{-3}$	–	$(3.3 \times 10^6)$
	$^6D_{9/2} - ^6D_{7/2}$	25.9882	$2.13 \times 10^{-3}$	–	$(2.2 \times 10^6)$

